

# Rac GAP1 Polyclonal Antibody

Catalog # AP72136

## **Product Information**

| Application       | WB, IHC-P, IF             |
|-------------------|---------------------------|
| Primary Accession | <u>Q9H0H5</u>             |
| Reactivity        | Human, Mouse, Rat, Monkey |
| Host              | Rabbit                    |
| Clonality         | Polyclonal                |
| Calculated MW     | 71027                     |

#### **Additional Information**

| Gene ID            | 29127  |
|--------------------|--|
| Other Names        | RACGAP1; KIAA1478; MGCRACGAP; Rac GTPase-activating protein 1; Male<br>germ cell RacGap; MgcRacGAP; Protein CYK4 homolg; CYK4; HsCYK-4   |
| Dilution           | WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.<br>Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other<br>applications. IHC-P~~N/A IF~~1:50~200 |
| Format             | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.  |
| Storage Conditions | -20°C  |

### **Protein Information**

| Name     | RACGAP1 ( <u>HGNC:9804</u> )  |
|----------|---|
| Function | Component of the centralspindlin complex that serves as a<br>microtubule-dependent and Rho-mediated signaling required for the myosin<br>contractile ring formation during the cell cycle cytokinesis. Required for<br>proper attachment of the midbody to the cell membrane during cytokinesis.<br>Sequentially binds to ECT2 and RAB11FIP3 which regulates cleavage furrow<br>ingression and abscission during cytokinesis (PubMed: <u>18511905</u> ). Plays key<br>roles in controlling cell growth and differentiation of hematopoietic cells<br>through mechanisms other than regulating Rac GTPase activity<br>(PubMed: <u>10979956</u> ). Has a critical role in erythropoiesis (PubMed: <u>34818416</u> ).<br>Also involved in the regulation of growth-related processes in adipocytes and<br>myoblasts. May be involved in regulating spermatogenesis and in the<br>RACGAP1 pathway in neuronal proliferation. Shows strong GAP (GTPase<br>activation) activity towards CDC42 and RAC1 and less towards RHOA. Essential<br>for the early stages of embryogenesis. May play a role in regulating cortical<br>activity through RHOA during cytokinesis. May participate in the regulation of<br>sulfate transport in male germ cells. |

| Cellular Location | Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle Cytoplasmic vesicle,<br>secretory vesicle, acrosome. Cleavage furrow Midbody, Midbody ring. Cell<br>membrane; Peripheral membrane protein; Cytoplasmic side.<br>Note=Colocalizes with RND2 in Golgi-derived proacrosomal vesicles and the<br>acrosome (By similarity). During interphase, localized to the nucleus and<br>cytoplasm along with microtubules, in anaphase, is redistributed to the<br>central spindle and, in telophase and cytokinesis, to the midbody ring, also<br>called Flemming body. Colocalizes with RHOA at the myosin contractile ring<br>during cytokinesis. Colocalizes with ECT2 to the mitotic spindles during<br>anaphase/metaphase, the cleavage furrow during telophase and at the<br>midbody at the end of cytokinesis. Colocalizes with Cdc42 to spindle<br>microtubules from prometaphase to telophase. |
|-------------------|---|
| Tissue Location   | Highly expressed in testis, thymus and placenta. Expressed at lower levels in<br>spleen and peripheral blood lymphocytes In testis, expression is restricted to<br>germ cells with the highest levels of expression found in spermatocytes.<br>Expression is regulated in a cell cycle-dependent manner and peaks during<br>G2/M phase  |

## Background

Component of the centralspindlin complex that serves as a microtubule-dependent and Rho-mediated signaling required for the myosin contractile ring formation during the cell cycle cytokinesis. Required for proper attachment of the midbody to the cell membrane during cytokinesis. Plays key roles in controlling cell growth and differentiation of hematopoietic cells through mechanisms other than regulating Rac GTPase activity. Also involved in the regulation of growth-related processes in adipocytes and myoblasts. May be involved in regulating spermatogenesis and in the RACGAP1 pathway in neuronal proliferation. Shows strong GAP (GTPase activation) activity towards CDC42 and RAC1 and less towards RHOA. Essential for the early stages of embryogenesis. May play a role in regulating cortical activity through RHOA during cytokinesis. May participate in the regulation of sulfate transport in male germ cells.

#### Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.