

PRAK (MAPKAPK5) Antibody (T182)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7216a

Product Information

Application	WB, E
Primary Accession	Q8IW41
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB13233
Calculated MW	54220
Antigen Region	160-189

Additional Information

Gene ID	8550
Other Names	MAP kinase-activated protein kinase 5, MAPK-activated protein kinase 5, MAPKAP kinase 5, MAPKAP-K5, MAPKAPK-5, MK-5, MK5, p38-regulated/activated protein kinase, PRAK, MAPKAPK5, PRAK
Target/Specificity	This PRAK(MAPKAPK5) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 160-189 amino acids from human PRAK(MAPKAPK5).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PRAK (MAPKAPK5) Antibody (T182) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MAPKAPK5
Synonyms	PRAK
Function	Tumor suppressor serine/threonine-protein kinase involved in mTORC1

signaling and post-transcriptional regulation. Phosphorylates FOXO3, ERK3/MAPK6, ERK4/MAPK4, HSP27/HSPB1, p53/TP53 and RHEB. Acts as a tumor suppressor by mediating Ras-induced senescence and phosphorylating p53/TP53. Involved in post-transcriptional regulation of MYC by mediating phosphorylation of FOXO3: phosphorylation of FOXO3 leads to promote nuclear localization of FOXO3, enabling expression of miR-34b and miR-34c, 2 post-transcriptional regulators of MYC that bind to the 3'UTR of MYC transcript and prevent MYC translation. Acts as a negative regulator of mTORC1 signaling by mediating phosphorylation and inhibition of RHEB. Part of the atypical MAPK signaling via its interaction with ERK3/MAPK6 or ERK4/MAPK4: the precise role of the complex formed with ERK3/MAPK6 or ERK4/MAPK4 is still unclear, but the complex follows a complex set of phosphorylation events: upon interaction with atypical MAPK (ERK3/MAPK6 or ERK4/MAPK4), ERK3/MAPK6 (or ERK4/MAPK4) is phosphorylated and then mediates phosphorylation and activation of MAPKAPK5, which in turn phosphorylates ERK3/MAPK6 (or ERK4/MAPK4). Mediates phosphorylation of HSP27/HSPB1 in response to PKA/PRKACA stimulation, inducing F-actin rearrangement.

Cellular Location

Cytoplasm. Nucleus. Note=Translocates to the cytoplasm following phosphorylation and activation. Interaction with ERK3/MAPK6 or ERK4/MAPK4 and phosphorylation at Thr-182, activates the protein kinase activity, followed by translocation to the cytoplasm Phosphorylation by PKA/PRKACA at Ser-115 also induces nuclear export

Tissue Location

Expressed ubiquitously.


Background

MAPKAPK5 is a member of the serine/threonine kinase family. In response to cellular stress and proinflammatory cytokines, this kinase is activated through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38-alpha, and MAPK11/p38-beta. In vitro, this kinase phosphorylates heat shock protein HSP27 at its physiologically relevant sites.

References


New, L., et al., EMBO J. 17(12):3372-3384 (1998).
 Ni, H., et al., Biochem. Biophys. Res. Commun. 243(2):492-496 (1998).
 Sudo T., Biochem. Biophys. Res. Commun. 269:521-525(2000).

Images



Anti-MAPKAPK5 Antibody (T182) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Western blot analysis of MAPKAPK5 (arrow) using rabbit



polyclonal MAPKAPK5 Antibody (T182) (Cat.#AP7216a).
293 cell lysates (2 ug/lane) either nontransfected (Lane 1)
or transiently transfected (Lane 2) with the MAPKAPK5
gene.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.