

Ran Polyclonal Antibody

Catalog # AP72181

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P62826
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24423

Additional Information

Gene ID	5901
Other Names	RAN; ARA24; OK/SW-cl.81; GTP-binding nuclear protein Ran; Androgen receptor-associated protein 24; GTPase Ran; Ras-like protein TC4; Ras-related nuclear protein
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	RAN
Synonyms	ARA24 {ECO:0000303 PubMed:10400640}
Function	GTPase involved in nucleocytoplasmic transport, participating both to the import and the export from the nucleus of proteins and RNAs (PubMed: 10400640 , PubMed: 17209048 , PubMed: 26272610 , PubMed: 27306458 , PubMed: 8276887 , PubMed: 8636225 , PubMed: 8692944 , PubMed: 8896452 , PubMed: 9351834 , PubMed: 9428644 , PubMed: 9822603). Switches between a cytoplasmic GDP- and a nuclear GTP-bound state by nucleotide exchange and GTP hydrolysis (PubMed: 11336674 , PubMed: 26272610 , PubMed: 29040603 , PubMed: 7819259 , PubMed: 8636225 , PubMed: 8692944 , PubMed: 8896452 , PubMed: 9351834 , PubMed: 9428644 , PubMed: 9822603). Nuclear import receptors such as importin beta bind their substrates only in the absence of GTP-bound RAN and release them upon direct interaction with GTP-bound RAN, while export receptors behave in the opposite way. Thereby, RAN controls cargo loading and release by transport

receptors in the proper compartment and ensures the directionality of the transport (PubMed:[8896452](#), PubMed:[9351834](#), PubMed:[9428644](#)). Interaction with RANBP1 induces a conformation change in the complex formed by XPO1 and RAN that triggers the release of the nuclear export signal of cargo proteins (PubMed:[20485264](#)). RAN (GTP-bound form) triggers microtubule assembly at mitotic chromosomes and is required for normal mitotic spindle assembly and chromosome segregation (PubMed:[10408446](#), PubMed:[29040603](#)). Required for normal progress through mitosis (PubMed:[12194828](#), PubMed:[29040603](#), PubMed:[8421051](#)). The complex with BIRC5/survivin plays a role in mitotic spindle formation by serving as a physical scaffold to help deliver the RAN effector molecule TPX2 to microtubules (PubMed:[18591255](#)). Acts as a negative regulator of the kinase activity of VRK1 and VRK2 (PubMed:[18617507](#)). Enhances AR-mediated transactivation. Transactivation decreases as the poly-Gln length within AR increases (PubMed:[10400640](#)).

Cellular Location

Nucleus. Nucleus envelope. Cytoplasm, cytosol Cytoplasm. Melanosome Note=Predominantly nuclear during interphase (PubMed:[10679025](#), PubMed:[12194828](#), PubMed:[8421051](#)). Becomes dispersed throughout the cytoplasm during mitosis (PubMed:[12194828](#), PubMed:[8421051](#)). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:[17081065](#)).

Tissue Location


Expressed in a variety of tissues.

Background

GTPase involved in nucleocytoplasmic transport, participating both to the import and the export from the nucleus of proteins and RNAs (PubMed:[10400640](#), PubMed:[8276887](#), PubMed:[8896452](#), PubMed:[8636225](#), PubMed:[8692944](#), PubMed:[9351834](#), PubMed:[9428644](#), PubMed:[9822603](#), PubMed:[26272610](#)). Switches between a cytoplasmic GDP- and a nuclear GTP-bound state by nucleotide exchange and GTP hydrolysis (PubMed:[7819259](#), PubMed:[8896452](#), PubMed:[8636225](#), PubMed:[8692944](#), PubMed:[9351834](#), PubMed:[9428644](#), PubMed:[9822603](#), PubMed:[29040603](#), PubMed:[11336674](#), PubMed:[26272610](#)). Nuclear import receptors such as importin beta bind their substrates only in the absence of GTP-bound RAN and release them upon direct interaction with GTP-bound RAN, while export receptors behave in the opposite way. Thereby, RAN controls cargo loading and release by transport receptors in the proper compartment and ensures the directionality of the transport (PubMed:[8896452](#), PubMed:[9351834](#), PubMed:[9428644](#)). Interaction with RANBP1 induces a conformation change in the complex formed by XPO1 and RAN that triggers the release of the nuclear export signal of cargo proteins (PubMed:[20485264](#)). RAN (GTP-bound form) triggers microtubule assembly at mitotic chromosomes and is required for normal mitotic spindle assembly and chromosome segregation (PubMed:[10408446](#), PubMed:[29040603](#)). Required for normal progress through mitosis (PubMed:[8421051](#), PubMed:[12194828](#), PubMed:[29040603](#)). The complex with BIRC5/survivin plays a role in mitotic spindle formation by serving as a physical scaffold to help deliver the RAN effector molecule TPX2 to microtubules (PubMed:[18591255](#)). Acts as a negative regulator of the kinase activity of VRK1 and VRK2 (PubMed:[18617507](#)). Enhances AR-mediated transactivation. Transactivation decreases as the poly-Gln length within AR increases (PubMed:[10400640](#)).

Images

Western Blot analysis of various cells using Ran Polyclonal Antibody diluted at 1 : 2000



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.