

Siah-1/2 Polyclonal Antibody

Catalog # AP72479

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q8IUQ4 , Q43255
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31123

Additional Information

Gene ID	6477
Other Names	SIAH1; HUMSIAH; E3 ubiquitin-protein ligase SIAH1; Seven in absentia homolog 1; Siah-1; Siah-1a; SIAH2; E3 ubiquitin-protein ligase SIAH2; Seven in absentia homolog 2; Siah-2; hSiah2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SIAH1
Synonyms	HUMSIAH
Function	E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:14506261, PubMed:14645235, PubMed:14654780, PubMed:15064394, PubMed:16085652, PubMed:19224863, PubMed:20508617, PubMed:22483617, PubMed:28546513, PubMed:32430360, PubMed:33591310, PubMed:9334332, PubMed:9858595). E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:14506261, PubMed:14645235, PubMed:14654780, PubMed:15064394, PubMed:16085652, PubMed:19224863, PubMed:20508617, PubMed:22483617, PubMed:9334332, PubMed:9858595). Mediates E3 ubiquitin ligase activity either through direct binding to substrates or by functioning as the essential RING domain subunit of larger

E3 complexes (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (ELL2, MYB, POU2AF1, PML and RBBP8), a cell surface receptor (DCC), the cell-surface receptor-type tyrosine kinase FLT3, the cytoplasmic signal transduction molecules (KLF10/TIEG1 and NUMB), an antiapoptotic protein (BAG1), a microtubule motor protein (KIF22), a protein involved in synaptic vesicle function in neurons (SYP), a structural protein (CTNNB1) and SNCAIP (PubMed:[10747903](#), PubMed:[11146551](#), PubMed:[11389839](#), PubMed:[11389840](#), PubMed:[11483517](#), PubMed:[11483518](#), PubMed:[11752454](#), PubMed:[12072443](#)). Confers constitutive instability to HIPK2 through proteasomal degradation (PubMed:[18536714](#), PubMed:[33591310](#)). It is thereby involved in many cellular processes such as apoptosis, tumor suppression, cell cycle, axon guidance, transcription regulation, spermatogenesis and TNF signaling (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Has some overlapping function with SIAH2 (PubMed:[14506261](#), PubMed:[14645235](#), PubMed:[14654780](#), PubMed:[15064394](#), PubMed:[16085652](#), PubMed:[19224863](#), PubMed:[20508617](#), PubMed:[22483617](#), PubMed:[9334332](#), PubMed:[9858595](#)). Induces apoptosis in cooperation with PEG3 (By similarity). Upon nitric oxid (NO) generation that follows apoptotic stimulation, interacts with S-nitrosylated GAPDH, mediating the translocation of GAPDH to the nucleus (By similarity). GAPDH acts as a stabilizer of SIAH1, facilitating the degradation of nuclear proteins (By similarity). Mediates ubiquitination and degradation of EGLN2 and EGLN3 in response to the unfolded protein response (UPR), leading to their degradation and subsequent stabilization of ATF4 (By similarity). Also part of the Wnt signaling pathway in which it mediates the Wnt-induced ubiquitin- mediated proteasomal degradation of AXIN1 (PubMed:[28546513](#), PubMed:[32430360](#)).


Cellular Location

Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Partially nuclear

Tissue Location


Widely expressed at a low level. Down-regulated in advanced hepatocellular carcinomas.

Images



Western Blot analysis of various cells using Siah-1/2
Polyclonal Antibody diluted at 1 : 500

Western Blot analysis of HeLa cells using Siah-1/2
Polyclonal Antibody diluted at 1 : 500



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.