

Skp2 p45 Polyclonal Antibody

Catalog # AP72494

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q13309
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47761

Additional Information

Gene ID	6502
Other Names	SKP2; FBXL1; S-phase kinase-associated protein 2; Cyclin-A/CDK2-associated protein p45; F-box protein Skp2; F-box/LRR-repeat protein 1; p45skp2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SKP2
Synonyms	FBXL1
Function	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed: 9736735 , PubMed: 11931757 , PubMed: 12435635 , PubMed: 12769844 , PubMed: 12840033 , PubMed: 15342634 , PubMed: 15668399 , PubMed: 15949444 , PubMed: 16103164 , PubMed: 16262255 , PubMed: 16581786 , PubMed: 16951159 , PubMed: 17908926 , PubMed: 17962192 , PubMed: 22464731 , PubMed: 22770219 , PubMed: 32267835). Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition (By similarity). Degradation of CDKN1B/p27kip also requires CKS1 (By similarity). Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, NBN, FOXO1, UBP43, YTHDF2, and probably MYC, TOB1 and TAL1 (PubMed: 11931757 , PubMed: 12435635 ,

PubMed:[12769844](#), PubMed:[12840033](#), PubMed:[15342634](#), PubMed:[15668399](#), PubMed:[15949444](#), PubMed:[16103164](#), PubMed:[16581786](#), PubMed:[16951159](#), PubMed:[17908926](#), PubMed:[17962192](#), PubMed:[22464731](#), PubMed:[32267835](#)). Degradation of TAL1 also requires STUB1 (PubMed:[17962192](#)). Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2 (PubMed:[9736735](#), PubMed:[16262255](#)). Promotes ubiquitination and destruction of CDH1 in a CK1-dependent manner, thereby regulating cell migration (PubMed:[22770219](#)). Following phosphorylation in response to DNA damage, mediates 'Lys-63'-linked ubiquitination of NBN, promoting ATM recruitment to DNA damage sites and DNA repair via homologous recombination (PubMed:[22464731](#)).


Cellular Location

Cytoplasm. Nucleus


Background

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription. Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition. Degradation of CDKN1B/p27kip also requires CKS1. Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, FOXO1, UBP43, and probably MYC, TOB1 and TAL1. Degradation of TAL1 also requires STUB1. Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2. Promotes ubiquitination and destruction of CDH1 in a CK1-Dependent Manner, thereby regulating cell migration.

Images





Western Blot analysis of various cells using Skp2 p45 Polyclonal Antibody diluted at 1 : 500



Immunohistochemical analysis of paraffin-embedded Human Right kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human Right kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human Right kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.