

# SLU7 Polyclonal Antibody

Catalog # AP72505

## Product Information

---

Application	WB
Primary Accession	<a href="#">O95391</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68387

## Additional Information

---

Gene ID	10569
Other Names	SLU7; Pre-mRNA-splicing factor SLU7; hSlu7
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	SLU7
Function	Required for pre-mRNA splicing as component of the spliceosome (PubMed: <a href="#">10197984</a> , PubMed: <a href="#">28502770</a> , PubMed: <a href="#">30705154</a> ). Participates in the second catalytic step of pre-mRNA splicing, when the free hydroxyl group of exon I attacks the 3'-splice site to generate spliced mRNA and the excised lariat intron. Required for holding exon 1 properly in the spliceosome and for correct AG identification when more than one possible AG exists in 3'-splicing site region. May be involved in the activation of proximal AG. Probably also involved in alternative splicing regulation.
Cellular Location	Nucleus. Nucleus speckle. Cytoplasm Note=Predominantly nuclear. Shuttling between the nucleus and the cytoplasm is regulated by the CCHC-type zinc finger. Upon UV-C stress stimulus, the nuclear concentration of the protein decreases, affecting alternative splicing. Translocates from the nucleus to the cytoplasm after heat shock cell treatment. Accumulates in cytoplasmic vesicle-like organelles after heat shock treatment, which may represent stress granules.


## Background

---

Participates in the second catalytic step of pre-mRNA splicing, when the free hydroxyl group of exon I attacks the 3'- splice site to generate spliced mRNA and the excised lariat intron. Required for holding exon 1 properly in the spliceosome and for correct AG identification when more than one possible AG exists in 3'-splicing site region. May be involved in the activation of proximal AG. Probably also involved in alternative splicing regulation.

## Images

---



Western Blot analysis of SH-SY5Y cells using SLU7  
Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.