

TADA2L Polyclonal Antibody

Catalog # AP72707

Product Information

Application WB Primary Accession 075478

Reactivity Human, Mouse, Rat, Monkey

HostRabbitClonalityPolyclonalCalculated MW51506

Additional Information

Gene ID 6871

Other Names TADA2A; TADA2L; KL04P; Transcriptional adapter 2-alpha; Transcriptional

adapter 2-like; ADA2-like protein

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name TADA2A

Synonyms TADA2L

Function Component of the ATAC complex, a complex with histone acetyltransferase

activity on histones H3 and H4. Required for the function of some acidic activation domains, which activate transcription from a distant site (By similarity). Binds double- stranded DNA. Binds dinucleosomes, probably at the linker region between neighboring nucleosomes. Plays a role in chromatin remodeling. May promote TP53/p53 'Lys-321' acetylation, leading to reduced TP53 stability and transcriptional activity (PubMed:22644376). May also promote XRCC6 acetylation thus facilitating cell apoptosis in response to DNA

damage (PubMed:22644376).


Cellular Location Nucleus. Chromosome {ECO:0000250 | UniProtKB:Q8CHV6}

Tissue Location Expressed in all tissues, but most abundantly in testis

Background

Component of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4. Required for the function of some acidic activation domains, which activate transcription from a distant site (By similarity). Binds double- stranded DNA. Binds dinucleosomes, probably at the linker region between neighboring nucleosomes. Plays a role in chromatin remodeling. May promote TP53/p53 'Lys-321' acetylation, leading to reduced TP53 stability and transcriptional activity (PubMed:22644376). May also promote XRCC6 acetylation thus facilitating cell apoptosis in response to DNA damage (PubMed:22644376).

Images



Western Blot analysis of various cells using TADA2L Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.