

REA (PHB2) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7270a

Product Information

Application	WB, IHC-P, E
Primary Accession	Q99623
Other Accession	Q5XIH7 , Q35129 , Q5ZMN3 , Q2HJ97 , NP_009204
Reactivity	Human
Predicted	Bovine, Chicken, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10629
Calculated MW	33296
Antigen Region	66-95

Additional Information

Gene ID	11331
Other Names	Prohibitin-2, B-cell receptor-associated protein BAP37, D-prohibitin, Repressor of estrogen receptor activity, PHB2 {ECO:0000312 EMBL:AAH147661, ECO:0000312 HGNC:HGNC:30306}
Target/Specificity	This REA (PHB2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 66-95 amino acids from the N-terminal region of human REA (PHB2).
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	REA (PHB2) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PHB2 {ECO:0000312 EMBL:AAH14766.1, ECO:0000312 HGNC:HGNC:30306}
Function	Protein with pleiotropic attributes mediated in a cell- compartment- and

tissue-specific manner, which include the plasma membrane-associated cell signaling functions, mitochondrial chaperone, and transcriptional co-regulator of transcription factors and sex steroid hormones in the nucleus.

Cellular Location

Mitochondrion inner membrane. Cytoplasm. Nucleus. Cell membrane
Note=Localizes within both nucleus and cytoplasm in proliferative primary myoblasts and mostly within the nucleus of differentiated primary myoblasts.
[Isoform 2]: Mitochondrion inner membrane

Tissue Location

Expressed in myoblasts.


Background

PHB2 acts as a mediator of transcriptional repression by nuclear hormone receptors via recruitment of histone deacetylases. This protein functions as an estrogen receptor (ER)-selective coregulator that potentiates the inhibitory activities of antiestrogens and represses the activity of estrogens. It competes with NCOA1 for modulation of ER transcriptional activity. It is probably involved in regulating mitochondrial respiration activity and in aging.


References

Takata,H., Curr. Biol. 17 (15), 1356-1361 (2007)
Kasashima,K., J. Biol. Chem. 281 (47), 36401-36410 (2006)

Images



Western blot analysis of PHB2 (arrow) using rabbit polyclonal PHB2 Antibody (Human C-term) (Cat.#AP7270a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PHB2 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with PHB2 Antibody (N-term) (Cat.#AP7270a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.