10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

REA (PHB2) Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7270b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q99623
Other Accession	Q5XIH7, O35129, Q2HJ97, NP_009204
Reactivity	Human
Predicted	Bovine, Mouse, Rat, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10631
Calculated MW	33296
Antigen Region	197-226

Additional Information

Gene ID	11331
Other Names	Prohibitin-2, B-cell receptor-associated protein BAP37, D-prohibitin, Repressor of estrogen receptor activity, PHB2 {ECO:0000312|EMBL:AAH147661, ECO:0000312|HGNC:HGNC:30306}
Target/Specificity	This REA (PHB2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 197-226 amino acids from the C-terminal region of human REA (PHB2).
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	REA (PHB2) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PHB2 {ECO:0000312|EMBL:AAH14766.1, ECO:0000312|HGNC:HGNC:30306}
Function	Protein with pleiotropic attributes mediated in a cell- compartment- and

	tissue-specific manner, which include the plasma membrane-associated cell signaling functions, mitochondrial chaperone, and transcriptional co-regulator of transcription factors and sex steroid hormones in the nucleus.
Cellular Location	Mitochondrion inner membrane. Cytoplasm. Nucleus. Cell membrane Note=Localizes within both nucleus and cytoplasm in proliferative primary myoblasts and mostly within the nucleus of differentiated primary myoblasts. [Isoform 2]: Mitochondrion inner membrane
Tissue Location	Expressed in myoblasts.


Background

Acts as a mediator of transcriptional repression by nuclear hormone receptors via recruitment of histone deacetylases. Functions as an estrogen receptor (ER)-selective coregulator that potentiates the inhibitory activities of antiestrogens and represses the activity of estrogens. Competes with NCOA1 for modulation of ER transcriptional activity. Probably involved in regulating mitochondrial respiration activity and in aging.


References

Takata,H., Curr. Biol. 17 (15), 1356-1361 (2007)
Kasashima,K., J. Biol. Chem. 281 (47), 36401-36410 (2006)

Images




All lanes : Anti-PHB2 (Human C-term) at 1:1000 dilution Lane 1: 293 whole cell lysate Lane 2: Hela whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of anti-PHB2 Antibody (C-term) (Cat. #AP7270b) in 293 cell line lysates (35ug/lane). PHB2(arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with PHB2 Antibody (C-term) (Cat.#AP7270b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been



evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.