

# TEF-4 Polyclonal Antibody

Catalog # AP72780

## **Product Information**

| Application       | WB, IHC-P, IF |
|-------------------|---------------|
| Primary Accession | <u>Q15562</u> |
| Reactivity        | Human, Mouse  |
| Host              | Rabbit        |
| Clonality         | Polyclonal    |
| Calculated MW     | 49243         |

#### **Additional Information**

| Gene ID            | 8463  |
|--------------------|---|
| Other Names        | TEAD2; TEF4; Transcriptional enhancer factor TEF-4; TEA domain family<br>member 2; TEAD-2 |
| Dilution           | WB~~1:1000 IHC-P~~N/A IF~~1:50~200  |
| Format             | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.             |
| Storage Conditions | -20°C   |

## **Protein Information**

| Name              | TEAD2  |
|-------------------|--|
| Synonyms          | TEF4   |
| Function          | Transcription factor which plays a key role in the Hippo signaling pathway, a pathway involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Acts by mediating gene expression of YAP1 and WWTR1/TAZ, thereby regulating cell proliferation, migration and epithelial mesenchymal transition (EMT) induction. Binds to the SPH and GT-IIC 'enhansons' (5'-GTGGAATGT-3'). May be involved in the gene regulation of neural development. Binds to the M-CAT motif. |
| Cellular Location | Nucleus.   |
| Background        |  |

Transcription factor which plays a key role in the Hippo signaling pathway, a pathway involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Acts by mediating gene expression of YAP1 and WWTR1/TAZ, thereby regulating cell proliferation, migration and epithelial mesenchymal transition (EMT) induction. Binds to the SPH and GT-IIC 'enhansons' (5'- GTGGAATGT-3'). May be involved in the gene regulation of neural development. Binds to the M-CAT motif.

### Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.