

# TFIID Polyclonal Antibody

Catalog # AP72798

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">P20226</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	37698

## Additional Information

---

<b>Gene ID</b>	6908
<b>Other Names</b>	TBP; GTF2D1; TF2D; TFIID; TATA-box-binding protein; TATA sequence-binding protein; TATA-binding factor; TATA-box factor; Transcription initiation factor TFIID TBP subunit
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	TBP
<b>Synonyms</b>	GTF2D1, TF2D, TFIID {ECO:0000303   PubMed:
<b>Function</b>	The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed: <a href="#">33795473</a> ). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed: <a href="#">2194289</a> , PubMed: <a href="#">2363050</a> , PubMed: <a href="#">2374612</a> , PubMed: <a href="#">27193682</a> , PubMed: <a href="#">33795473</a> ). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed: <a href="#">27007846</a> , PubMed: <a href="#">33795473</a> ). The TFIID complex structure can be divided into 3 modules TFIID-A, TFIID-B, and TFIID-C (PubMed: <a href="#">33795473</a> ). TBP forms the TFIID-A module together with TAF3 and

TAF5 (PubMed:[33795473](#)). TBP is a general transcription factor that functions at the core of the TFIID complex (PubMed:[2194289](#), PubMed:[2363050](#), PubMed:[2374612](#), PubMed:[27193682](#), PubMed:[33795473](#), PubMed:[9836642](#)). During assembly of the core PIC on the promoter, as part of TFIID, TBP binds to and also bends promoter DNA, irrespective of whether the promoter contains a TATA box (PubMed:[33795473](#)). Component of a BRF2-containing transcription factor complex that regulates transcription mediated by RNA polymerase III (PubMed:[26638071](#)). Component of the transcription factor SL1/TIF-IB complex, which is involved in the assembly of the PIC during RNA polymerase I-dependent transcription (PubMed:[15970593](#)). The rate of PIC formation probably is primarily dependent on the rate of association of SL1 with the rDNA promoter (PubMed:[15970593](#)). SL1 is involved in stabilization of nucleolar transcription factor 1/UBTF on rDNA (PubMed:[15970593](#)).

#### Cellular Location

Nucleus.

#### Tissue Location

Widely expressed, with levels highest in the testis and ovary.


## Background

General transcription factor that functions at the core of the DNA-binding multiprotein factor TFIID (PubMed:[2374612](#), PubMed:[2363050](#), PubMed:[2194289](#), PubMed:[9836642](#), PubMed:[27193682](#)). Binding of TFIID to the TATA box is the initial transcriptional step of the pre-initiation complex (PIC), playing a role in the activation of eukaryotic genes transcribed by RNA polymerase II (PubMed:[2374612](#), PubMed:[2363050](#), PubMed:[2194289](#), PubMed:[9836642](#), PubMed:[27193682](#)). Component of a BRF2-containing transcription factor complex that regulates transcription mediated by RNA polymerase III (PubMed:[26638071](#)). Component of the transcription factor SL1/TIF-IB complex, which is involved in the assembly of the PIC (pre-initiation complex) during RNA polymerase I-dependent transcription (PubMed:[15970593](#)). The rate of PIC formation probably is primarily dependent on the rate of association of SL1 with the rDNA promoter. SL1 is involved in stabilization of nucleolar transcription factor 1/UBTF on rDNA.

## Images



Western Blot analysis of various cells using TFIID Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.