

TGFβ2 Polyclonal Antibody

Catalog # AP72816

Product Information

Application WB, IHC-P **Primary Accession** P61812

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW47748

Additional Information

Gene ID 7042

Other Names TGFB2; Transforming growth factor beta-2; TGF-beta-2; BSC-1 cell growth

inhibitor; Cetermin; Glioblastoma-derived T-cell suppressor factor; G-TSF;

Polyergin

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name TGFB2

Function [Transforming growth factor beta-2 proprotein]: Precursor of the

Latency-associated peptide (LAP) and Transforming growth factor beta-2 (TGF-beta-2) chains, which constitute the regulatory and active subunit of

TGF-beta-2, respectively.


Cellular Location [Latency-associated peptide]: Secreted, extracellular space, extracellular

matrix {ECO:0000250 | UniProtKB:P01137}

Background

Transforming growth factor beta-2 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-2 (TGF-beta-2) chains, which constitute the regulatory and active subunit of TGF-beta-2, respectively.

Images



Western Blot analysis of various cells using TGF β 2 Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.