

# Tra-2 $\alpha$ Polyclonal Antibody

Catalog # AP72896

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q13595</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	32689

## Additional Information

---

<b>Gene ID</b>	29896
<b>Other Names</b>	TRA2A; Transformer-2 protein homolog alpha; TRA-2 alpha; TRA2-alpha; Transformer-2 protein homolog A
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	TRA2A ( <a href="#">HGNC:16645</a> )
<b>Function</b>	Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.
<b>Cellular Location</b>	Nucleus.

## Background


---

Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.

## Images

---

Western Blot analysis of various cells using Tra-2 $\alpha$  Polyclonal Antibody diluted at 1 : 500. Secondary



antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.