10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TTF-I Polyclonal Antibody

Catalog # AP72954

Product Information

Application	IHC-P
Primary Accession	Q15361
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	103051

Additional Information

Gene ID	7270
Other Names	TTF1; Transcription termination factor 1; TTF-1; RNA polymerase I termination factor; Transcription termination factor I; TTF-I
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information

Name	TTF1
Function	Multifunctional nucleolar protein that terminates ribosomal gene transcription, mediates replication fork arrest and regulates RNA polymerase I transcription on chromatin. Plays a dual role in rDNA regulation, being involved in both activation and silencing of rDNA transcription. Interaction with BAZ2A/TIP5 recovers DNA-binding activity.
Cellular Location	Nucleus {ECO:0000250|UniProtKB:Q62187}. Nucleus, nucleolus {ECO:0000250|UniProtKB:Q62187}. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q62187}. Note=May be localized to the nucleolus in an NPM1/B23-dependent manner. May be displaced from the nucleolus into the nucleoplasm in an CDKN2A/ARF-dependent manner. May shuttle back and forth from nucleoplasm to nucleolus {ECO:0000250|UniProtKB:Q62187}

Background

Multifunctional nucleolar protein that terminates ribosomal gene transcription, mediates replication fork arrest and regulates RNA polymerase I transcription on chromatin. Plays a dual role in rDNA regulation,

being involved in both activation and silencing of rDNA transcription. Interaction with BAZ2A/TIP5 recovers DNA-binding activity.

Images



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.