

UBCE7IP4 Polyclonal Antibody

Catalog # AP72979

Product Information

Application	WB
Primary Accession	P50876
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32890

Additional Information

Gene ID	9781
Other Names	RNF144A; KIAA0161; RNF144; UBCE7IP4; Probable E3 ubiquitin-protein ligase RNF144A; RING finger protein 144A; UbcM4-interacting protein 4; Ubiquitin-conjugating enzyme 7-interacting protein 4
Dilution	WB~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	RNF144A
Synonyms	KIAA0161, RNF144, UBCE7IP4
Function	E3 ubiquitin-protein ligase which accepts ubiquitin from E2 ubiquitin-conjugating enzymes UBE2L3 and UBE2L6 in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed: 26216882). Mediates the ubiquitination and degradation of the DNA damage kinase PRKDC during DNA damage (PubMed: 24979766). Positively regulates DNA virus or exogenous cytosolic DNA-triggered innate immune response by mediating STING1 ubiquitination and increasing its 'Lys-6'-linked ubiquitination and translocation from the endoplasmic reticulum to the Golgi leading to downstream signaling pathways (PubMed: 37955227). Plays a positive role in EGF-dependent cell proliferation by prolonging EGF/EGFR signaling during EGF stimulation through EGFR ubiquitination (PubMed: 30171075). Increases ERK activity independently of EGFR signaling by promoting polyubiquitination and subsequent degradation of VRK3 in the cytosol (PubMed: 33067254).

Cellular Location

Cell membrane; Single-pass membrane protein. Cytoplasmic vesicle membrane. Endosome membrane. Endoplasmic reticulum membrane

Background

E3 ubiquitin-protein ligase which accepts ubiquitin from E2 ubiquitin-conjugating enzymes UBE2L3 and UBE2L6 in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Mediates the ubiquitination and degradation of the DNA damage kinase PRKDC.

Images



Western Blot analysis of various cells using UBCE7IP4 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.