

UMP-CMP Kinase Polyclonal Antibody

Catalog # AP73004

Product Information

Application	WB, IHC-P, IF
Primary Accession	P30085
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22222

Additional Information

Gene ID	51727
Other Names	CMPK1; CMK; CMPK; UCK; UMK; UMPK; UMP-CMP kinase; Cytidine monophosphate kinase; Cytidylate kinase; Deoxycytidylate kinase; Uridine monophosphate kinase; Uridine monophosphate/cytidine monophosphate kinase; UMP/CMP kinase; UMP/CMPK
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CMPK1 {ECO:0000255 HAMAP-Rule:MF_03172}
Function	Catalyzes the phosphorylation of pyrimidine nucleoside monophosphates at the expense of ATP. Plays an important role in de novo pyrimidine nucleotide biosynthesis. Has preference for UMP and CMP as phosphate acceptors. Also displays broad nucleoside diphosphate kinase activity.
Cellular Location	Nucleus {ECO:0000255 HAMAP-Rule:MF_03172, ECO:0000269 PubMed:10462544, ECO:0000269 PubMed:11912132}. Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03172, ECO:0000269 PubMed:10462544, ECO:0000269 PubMed:11912132}. Note=Predominantly cytoplasmic, less than 15% nuclear.
Tissue Location	Ubiquitously expressed.

Background

Catalyzes the phosphorylation of pyrimidine nucleoside monophosphates at the expense of ATP. Plays an important role in de novo pyrimidine nucleotide biosynthesis. Has preference for UMP and CMP as phosphate acceptors. Also displays broad nucleoside diphosphate kinase activity.

Images



Western Blot analysis of various cells using UMP-CMP Kinase Polyclonal Antibody diluted at 1 : 500. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.