

USP38 Polyclonal Antibody

Catalog # AP73020

Product Information

Application	WB, E
Primary Accession	Q8NB14
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116546

Additional Information

Gene ID	84640
Other Names	USP38; KIAA1891; Ubiquitin carboxyl-terminal hydrolase 38; Deubiquitinating enzyme 38; HP43.8KD; Ubiquitin thioesterase 38; Ubiquitin-specific-processing protease 38
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	USP38
Synonyms	KIAA1891
Function	Deubiquitinating enzyme that plays a role in various cellular processes, including DNA repair, cell cycle regulation, and immune response (PubMed:22689415, PubMed:30497519, PubMed:31874856, PubMed:35238669). Plays a role in the inhibition of type I interferon signaling by mediating the 'Lys-33' to 'Lys-48' ubiquitination transition of TBK1 leading to its degradation (PubMed:27692986). Cleaves the ubiquitin chain from the histone demethylase LSD1/KDM1A and prevents it from degradation by the 26S proteasome, thus maintaining LSD1 protein level in cells (PubMed:30497519). Plays a role in the DNA damage response by regulating the deacetylase activity of HDAC1 (PubMed:31874856). Mechanistically, removes the 'Lys-63'-linked ubiquitin chain promoting the deacetylase activity of HDAC1 in response to DNA damage (PubMed:31874856). Also acts as a specific deubiquitinase of histone deacetylase 3/HDAC3 and cleaves its 'Lys-63'-linked ubiquitin chains to lower its histone deacetylase activity

(PubMed:[32404892](#)). Regulates MYC levels and cell proliferation via antagonizing ubiquitin E3 ligase FBXW7 thereby preventing MYC 'Lys-48'-linked ubiquitination and degradation (PubMed:[34102342](#)). Participates in antiviral response by removing both 'Lys-48'-linked and 'Lys-63'-linked polyubiquitination of Zika virus envelope protein E (PubMed:[34696459](#)). Constitutively associated with IL-33R/IL1RL1, deconjugates its 'Lys-27'-linked polyubiquitination resulting in its autophagic degradation (PubMed:[35238669](#)).

Cellular Location

Cytoplasm. Nucleus Note=In response to DNA damage, recruited to DNA damage sites in the nucleus.


Tissue Location

Highly expressed in skeletal muscle. Expressed in adrenal gland.

Background

Deubiquitinating enzyme exhibiting a preference towards 'Lys-63'-linked ubiquitin chains.

Images



Western Blot analysis of various cells using USP38 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.