

Vasohibin Polyclonal Antibody

Catalog # AP73036

Product Information

Application	WB, E
Primary Accession	Q7L8A9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40957

Additional Information

Gene ID	22846
Other Names	VASH1; KIAA1036; VASH; Vasohibin-1
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	VASH1 (HGNC:19964)
Function	Tyrosine carboxypeptidase that removes the C-terminal tyrosine residue of alpha-tubulin, thereby regulating microtubule dynamics and function (PubMed:29146869, PubMed:31171830, PubMed:31235910, PubMed:31235911, PubMed:31270470). Critical for spindle function and accurate chromosome segregation during mitosis since microtubule detyrosination regulates mitotic spindle length and positioning (PubMed:31171830). Acts as an angiogenesis inhibitor: inhibits migration, proliferation and network formation by endothelial cells as well as angiogenesis (PubMed:15467828, PubMed:16488400, PubMed:16707096, PubMed:19204325). This inhibitory effect is selective to endothelial cells as it does not affect the migration of smooth muscle cells or fibroblasts (PubMed:15467828, PubMed:16488400, PubMed:16707096).
Cellular Location	Cytoplasm. Secreted. Note=Mainly localizes in the cytoplasm (PubMed: 27879017). Some fraction is secreted via a non-canonical secretion system; interaction with SVBP promotes secretion (PubMed: 27879017).
Tissue Location	Preferentially expressed in endothelial cells (PubMed: 15467828 ,

PubMed:16707096). Highly expressed in fetal organs (PubMed:15467828). Expressed in brain and placenta, and at lower level in heart and kidney (PubMed:15467828). Highly detected in microvessels endothelial cells of atherosclerotic lesions (PubMed:16707096)

Background

Tyrosine carboxypeptidase that removes the C-terminal tyrosine residue of alpha-tubulin, thereby regulating microtubule dynamics and function (PubMed:[29146869](#)). Acts as an angiogenesis inhibitor: inhibits migration, proliferation and network formation by endothelial cells as well as angiogenesis (PubMed:[15467828](#), PubMed:[16488400](#), PubMed:[16707096](#), PubMed:[19204325](#)). This inhibitory effect is selective to endothelial cells as it does not affect the migration of smooth muscle cells or fibroblasts (PubMed:[15467828](#), PubMed:[16488400](#), PubMed:[16707096](#)).

Images



Western Blot analysis of various cells using Vasohibin Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.