

VEGF-B Polyclonal Antibody

Catalog # AP73054

Product Information

Application	WB, IHC-P
Primary Accession	P49765
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21602

Additional Information

Gene ID	7423
Other Names	VEGFB; VRF; Vascular endothelial growth factor B; VEGF-B; VEGF-related factor; VRF
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	VEGFB
Synonyms	VRF
Function	Growth factor for endothelial cells. VEGF-B167 binds heparin and neuropilin-1 whereas the binding to neuropilin-1 of VEGF-B186 is regulated by proteolysis.
Cellular Location	Secreted. Note=Secreted but remains associated to cells or to the extracellular matrix unless released by heparin
Tissue Location	Expressed in all tissues except liver. Highest levels found in heart, skeletal muscle and pancreas

Background

Growth factor for endothelial cells. VEGF-B167 binds heparin and neuropilin-1 whereas the binding to neuropilin-1 of VEGF-B186 is regulated by proteolysis.

Images



Western Blot analysis of various cells using VEGF-B
Polyclonal Antibody diluted at 1 : 2000. Secondary
antibody was diluted at 1:20000



Western Blot analysis of Jurkat cells using VEGF-B
Polyclonal Antibody diluted at 1 : 2000. Secondary
antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.