10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Versican Polyclonal Antibody

Catalog # AP73057

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P13611
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	372820

Additional Information

Gene ID	1462
Other Names	VCAN; CSPG2; Versican core protein; Chondroitin sulfate proteoglycan core protein 2; Chondroitin sulfate proteoglycan 2; Glial hyaluronate-binding protein; GHAP; Large fibroblast proteoglycan; PG-M
Dilution	WB~~Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	VCAN
Synonyms	CSPG2
Function	May play a role in intercellular signaling and in connecting cells with the extracellular matrix. May take part in the regulation of cell motility, growth and differentiation. Binds hyaluronic acid.
Cellular Location	Secreted, extracellular space, extracellular matrix. Cell projection, cilium, photoreceptor outer segment. Secreted, extracellular space, extracellular matrix, interphotoreceptor matrix. Secreted
Tissue Location	Detected in placenta (at protein level) (PubMed:32337544). Detected in cerebrospinal fluid, fibroblasts and urine (at protein level) (PubMed:25326458, PubMed:36213313). Expressed in the retina (at protein level) (PubMed:29777959). Cerebral white matter and plasma (PubMed:2469524). Isoform V0: Expressed in normal brain, gliomas, medulloblastomas, schwannomas, neurofibromas, and meningiomas

(PubMed:8627343). Isoform V1: Expressed in normal brain, gliomas, medulloblastomas, schwannomas, neurofibromas, and meningiomas (PubMed:8627343). Isoform V2: Restricted to normal brain and gliomas (PubMed:8627343). Isoform V3: Found in all these tissues except medulloblastomas (PubMed:8627343).

Background

May play a role in intercellular signaling and in connecting cells with the extracellular matrix. May take part in the regulation of cell motility, growth and differentiation. Binds hyaluronic acid.

Images



Western Blot analysis of mouse-brain cells using Versican Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.