

Vinculin Polyclonal Antibody

Catalog # AP73065

Product Information

Application	WB, IHC-P, IF
Primary Accession	P18206
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	123799

Additional Information

Gene ID	7414
Other Names	VCL; Vinculin; Metavinculin
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	VCL
Function	Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250 UniProtKB:P12003}; Cytoplasmic side {ECO:0000250 UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250 UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250 UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250 UniProtKB:Q64727}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250 UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250 UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions

Tissue Location Metavinculin is muscle-specific.

Background

Actin filament (F-actin)-binding protein involved in cell-matrix adhesion and cell-cell adhesion. Regulates cell- surface E-cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.

Images



Western Blot analysis of various cells using Vinculin Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Citations

- [Sustained adenosine exposure causes endothelial mitochondrial dysfunction via equilibrative nucleoside transporters](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.