

VIP Polyclonal Antibody

Catalog # AP73066

Product Information

Application	IHC-P, IF, ICC, WB, E
Primary Accession	P01282
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19169

Additional Information

Gene ID	7432
Other Names	VIP; VIP peptides
Dilution	IHC-P~~N/A IF~~1:50~200 ICC~~N/A WB~~WB 1:500-2000 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	VIP (HGNC:12693)
Function	[Vasoactive intestinal peptide]: VIP is a neuropeptide involved in a diverse array of physiological processes through activating the PACAP subfamily of class B1 G protein-coupled receptors: VIP receptor 1 (VPR1) and VIP receptor 2 (VPR2) (PubMed: 1318039 , PubMed: 36385145 , PubMed: 8933357). Abundantly expressed throughout the CNS and peripheral nervous systems where they primarily exert neuroprotective and immune modulatory roles (PubMed: 3456568). Also causes vasodilation, lowers arterial blood pressure, stimulates myocardial contractility, increases glycogenolysis and relaxes the smooth muscle of trachea, stomach and gall bladder (PubMed: 15013843).
Cellular Location	Secreted.

Background

VIP causes vasodilation, lowers arterial blood pressure, stimulates myocardial contractility, increases glycogenolysis and relaxes the smooth muscle of trachea, stomach and gall bladder.

Images



Immunohistochemical analysis of paraffin-embedded Human testis. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.