

Wnt-1 Polyclonal Antibody

Catalog # AP73090

Product Information

Application	WB, IHC-P, IF
Primary Accession	P04628
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40982

Additional Information

Gene ID	7471
Other Names	WNT1; INT1; Proto-oncogene Wnt-1; Proto-oncogene Int-1 homolog
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	WNT1
Synonyms	INT1
Function	Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Acts in the canonical Wnt signaling pathway by promoting beta-catenin-dependent transcriptional activation (PubMed: 23499309 , PubMed: 23656646 , PubMed: 26902720 , PubMed: 28528193). In some developmental processes, is also a ligand for the coreceptor RYK, thus triggering Wnt signaling (By similarity). Plays an essential role in the development of the embryonic brain and central nervous system (CNS) (By similarity). Has a role in osteoblast function, bone development and bone homeostasis (PubMed: 23499309 , PubMed: 23656646).
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted

Background

Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Acts in the canonical Wnt signaling pathway by promoting beta-catenin-dependent transcriptional activation (PubMed:[23499309](#), PubMed:[26902720](#), PubMed:[28528193](#), PubMed:[23656646](#)). In some developmental processes, is also a ligand for the coreceptor RYK, thus triggering Wnt signaling (By similarity). Plays an essential role in the development of the embryonic brain and central nervous system (CNS) (By similarity). Has a role in osteoblast function, bone development and bone homeostasis (PubMed:[23499309](#), PubMed:[23656646](#)).

Images



Western Blot analysis of various cells using Wnt-1 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000



Western Blot analysis of HT29 cells using Wnt-1 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.