

WTAP Polyclonal Antibody

Catalog # AP73091

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q15007
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44244

Additional Information

Gene ID	9589
Other Names	WTAP; KIAA0105; Pre-mRNA-splicing regulator WTAP; Female-lethal(2)D homolog; hFL(2)D; WT1-associated protein; Wilms tumor 1-associating protein
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	WTAP {ECO:0000303 PubMed:11001926, ECO:0000312 HGNC:HGNC:16846}
Function	Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed: 29507755). Required for accumulation of METTL3 and METTL14 to nuclear speckle (PubMed: 24316715 , PubMed: 24407421 , PubMed: 24981863). Acts as a mRNA splicing regulator (PubMed: 12444081). Regulates G2/M cell-cycle transition by binding to the 3' UTR of CCNA2, which enhances its stability (PubMed: 17088532). Impairs WT1 DNA-binding ability and inhibits expression of WT1 target genes (PubMed: 17095724).
Cellular Location	Nucleus speckle. Nucleus, nucleoplasm. Cytoplasm {ECO:0000250 UniProtKB:Q9ER69}. Note=Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the MACOM subcomplex, such as VIRMA, in the nucleus {ECO:0000250 UniProtKB:Q9ER69}


Tissue Location

Ubiquitously expressed.

Background

Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed:[29507755](#)). Required for accumulation of METTL3 and METTL14 to nuclear speckle (PubMed:[24316715](#), PubMed:[24407421](#), PubMed:[24981863](#)). Acts as a mRNA splicing regulator (PubMed:[12444081](#)). Regulates G2/M cell-cycle transition by binding to the 3' UTR of CCNA2, which enhances its stability (PubMed:[17088532](#)). Impairs WT1 DNA-binding ability and inhibits expression of WT1 target genes (PubMed:[17095724](#)).

Images



Western Blot analysis of various cells using WTAP Polyclonal Antibody diluted at 1 : 500. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.