

ZIP4 Polyclonal Antibody

Catalog # AP73132

Product Information

Application	WB, E
Primary Accession	Q6P5W5
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68408

Additional Information

Gene ID	55630
Other Names	SLC39A4; ZIP4; Zinc transporter ZIP4; Solute carrier family 39 member 4; Zrt- and Irt-like protein 4; ZIP-4
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SLC39A4 (HGNC:17129)
Synonyms	ZIP4
Function	Selective transporter that mediates the uptake of Zn(2+) (PubMed: 17202136 , PubMed: 22242765 , PubMed: 27321477 , PubMed: 28875161 , PubMed: 31164399 , PubMed: 31914589 , PubMed: 31979155 , PubMed: 33837739 , PubMed: 36473915). Plays an essential role for dietary zinc uptake from small intestine (By similarity). The Zn(2+) uniporter activity is regulated by zinc availability (PubMed: 17202136 , PubMed: 32348750). Also exhibits polyspecific binding and transport of Cu(2+), Cd(2+) and possibly Ni(2+) but at higher concentrations (PubMed: 22242765 , PubMed: 31914589).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:A0A0H3LM39}. Recycling endosome membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:A0A0H3LM39}. Apical cell membrane {ECO:0000250 UniProtKB:Q78IQ7}; Multi-pass membrane protein {ECO:0000250 UniProtKB:A0A0H3LM39}. Note=Colocalized with TFRC in the recycling endosomes. Cycles between endosomal compartments and

the plasma membrane in response to Zn(2+) availability. Zn(2+) deficiency promotes accumulation of SLC39A4 on the surface membrane, whereas high extracellular Zn(2+) levels induce internalization of SLC39A4, but also trigger drastic removal of cellular SLC39A4 via proteasomal and lysosomal degradation pathways. {ECO:0000250|UniProtKB:Q78IQ7, ECO:0000269|PubMed:17202136, ECO:0000269|PubMed:32348750}


Tissue Location

Highly expressed in kidney, small intestine, stomach, colon, jejunum and duodenum.

Background

Plays an important role in cellular zinc homeostasis as a zinc transporter. Regulated in response to zinc availability (By similarity).

Images



Western Blot analysis of Colo cells using ZIP4 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.