

ZIP7 Polyclonal Antibody

Catalog # AP73133

Product Information

Application	WB, E
Primary Accession	Q92504
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50118

Additional Information

Gene ID	7922
Other Names	SLC39A7; HKE4; RING5; Zinc transporter SLC39A7; Histidine-rich membrane protein Ke4; Really interesting new gene 5 protein; Solute carrier family 39 member 7
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	SLC39A7
Synonyms	HKE4 {ECO:0000303 PubMed:14525538}, RING
Function	Transports Zn(2+) from the endoplasmic reticulum (ER)/Golgi apparatus to the cytosol, playing an essential role in the regulation of cytosolic zinc levels (PubMed: 14525538 , PubMed: 15705588 , PubMed: 28205653 , PubMed: 29980658). Acts as a gatekeeper of zinc release from intracellular stores, requiring post-translational activation by phosphorylation, resulting in activation of multiple downstream pathways leading to cell growth and proliferation (PubMed: 22317921 , PubMed: 28205653 , PubMed: 29980658). Has an essential role in B cell development and is required for proper B cell receptor signaling (PubMed: 30718914). Plays an important role in maintaining intestinal epithelial homeostasis and skin dermis development by regulating ER function (By similarity). Controls cell signaling pathways involved in glucose metabolism in skeletal muscle (By similarity). Has a protective role against ER stress in different biological contexts (PubMed: 29980658 , PubMed: 30237509). Mediates Zn(2+)-induced ferroptosis (PubMed: 33608508).

Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus, cis-Golgi network membrane; Multi-pass membrane protein
Tissue Location	Widely expressed.

Background

Zinc transporter, that transports Zn(2+) from the endoplasmic reticulum/Golgi apparatus to the cytosol. Transport is stimulated by growth factors, such as EGF, and Ca(2+), as well as by exogenous Zn(2+).

Images



Western Blot analysis of various cells using ZIP7 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.