

ZnT-9 Polyclonal Antibody

Catalog # AP73174

Product Information

Application WB
Primary Accession Q6PML9

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW63515

Additional Information

Gene ID 10463

Other Names SLC30A9; C4orf1; HUEL; Zinc transporter 9; ZnT-9; Human embryonic lung

protein; HuEL; Solute carrier family 30 member 9

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name SLC30A9 (HGNC:1329)

Synonyms C4orf1, HUEL

Function Mitochondrial proton-coupled zinc ion antiporter mediating the export of

zinc from the mitochondria and involved in zinc homeostasis, zinc mobilization as well as mitochondrial morphology and health (PubMed: <u>28334855</u>, PubMed: <u>34433664</u>,

PubMed:<u>35614220</u>). In nucleus, functions as a secondary coactivator for nuclear receptors by cooperating with p160 coactivators subtypes. Plays a role in transcriptional activation of Wnt-responsive genes (By similarity).

Cellular Location Mitochondrion membrane; Multi-pass membrane protein. Nucleus

Endoplasmic reticulum. Note=Partial co- localization with endoplasmic reticulum (PubMed:28334855). Linked to mitochondrial ribosomes


(PubMed:35614220)

Tissue Location Ubiquitously expressed in fetal and adult tissues and cancer cell lines.

Background

Acts as a zinc transporter involved in intracellular zinc homeostasis (PubMed: <u>28334855</u>). Functions as a secondary coactivator for nuclear receptors by cooperating with p160 coactivators subtypes. Plays a role in transcriptional activation of Wnt-responsive genes (By similarity).

Images



Western Blot analysis of HepG2 cells using ZnT-9 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.