

α -protein Kinase 1 Polyclonal Antibody

Catalog # AP73184

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q96QP1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	138861

Additional Information

Gene ID	80216
Other Names	ALPK1; KIAA1527; LAK; Alpha-protein kinase 1; Chromosome 4 kinase; Lymphocyte alpha-protein kinase
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	ALPK1 {ECO:0000303 PubMed:30111836, ECO:0000312 HGNC:HGNC:20917}
Function	Serine/threonine-protein kinase that detects bacterial pathogen-associated molecular pattern metabolites (PAMPs) and initiates an innate immune response, a critical step for pathogen elimination and engagement of adaptive immunity (PubMed: 28222186 , PubMed: 28877472 , PubMed: 30111836). Specifically recognizes and binds ADP-D-glycero-beta- D-manno-heptose (ADP-Heptose), a potent PAMP present in all Gram- negative and some Gram-positive bacteria (PubMed: 30111836). ADP- Heptose-binding stimulates its kinase activity to phosphorylate and activate TIFA, triggering pro-inflammatory NF-kappa-B signaling (PubMed: 30111836 , PubMed: 35868845 , PubMed: 38060563). May be involved in monosodium urate monohydrate (MSU)-induced inflammation by mediating phosphorylation of unconventional myosin MYO9A (PubMed: 27169898). May also play a role in apical protein transport by mediating phosphorylation of unconventional myosin MYO1A (PubMed: 15883161). May play a role in ciliogenesis (PubMed: 30967659).

Cellular Location	Cytoplasm, cytosol. Cytoplasm, cytoskeleton, spindle pole Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium. Note=Localized at the base of primary cilia.
Tissue Location	Highly expressed in liver. Expressed in the optic nerve and retinal pigmented epithelium. Lower expression is observed in the macula and extramacular retina (PubMed:30967659)

Background

Serine/threonine-protein kinase that detects bacterial pathogen-associated molecular pattern metabolites (PAMPs) and initiates an innate immune response, a critical step for pathogen elimination and engagement of adaptive immunity (PubMed:[28877472](#), PubMed:[28222186](#), PubMed:[30111836](#)). Specifically recognizes and binds ADP-D-glycero-beta-D-manno-heptose (ADP-Heptose), a potent PAMP present in all Gram-negative and some Gram-positive bacteria (PubMed:[30111836](#)). ADP-Heptose-binding stimulates its kinase activity to phosphorylate and activate TIFA, triggering proinflammatory NF-kappa-B signaling (PubMed:[30111836](#)). May be involved in monosodium urate monohydrate (MSU)-induced inflammation by mediating phosphorylation of unconventional myosin MYO9A (PubMed:[27169898](#)). May also play a role in apical protein transport by mediating phosphorylation of unconventional myosin MYO1A (PubMed:[15883161](#)).

Images



Western Blot analysis of various cells using α-protein Kinase 1 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.