

EpoR Polyclonal Antibody

Catalog # AP73199

Product Information

Application	WB
Primary Accession	P19235
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55065

Additional Information

Gene ID	2057
Other Names	EPOR; Erythropoietin receptor; EPO-R
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	EPOR {ECO:0000303 PubMed:2163695, ECO:0000312 HGNC:HGNC:3416}
Function	Receptor for erythropoietin, which mediates erythropoietin- induced erythroblast proliferation and differentiation (PubMed: 10388848 , PubMed: 2163695 , PubMed: 2163696 , PubMed: 8662939 , PubMed: 9774108). Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade (By similarity). In some cell types, can also activate STAT1 and STAT3 (PubMed: 11756159). May also activate the LYN tyrosine kinase (By similarity).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P14753}; Single-pass type I membrane protein
Tissue Location	Erythroid cells and erythroid progenitor cells. [Isoform EPOR-S]: Isoform EPOR-S and isoform EPOR-T are the predominant forms in bone marrow.

Background

Receptor for erythropoietin. Mediates erythropoietin- induced erythroblast proliferation and differentiation. Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade. In some

cell types, can also activate STAT1 and STAT3. May also activate the LYN tyrosine kinase.

Images



Western Blot analysis of JK cells using EpoR Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.