

CHD1L Polyclonal Antibody

Catalog # AP73210

Product Information

Application	WB, E
Primary Accession	Q86WJ1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101000

Additional Information

Gene ID	9557
Other Names	CHD1L; ALC1; Chromodomain-helicase-DNA-binding protein 1-like; Amplified in liver cancer protein 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CHD1L {ECO:0000303 PubMed:34210977, ECO:0000312 HGNC:HGNC:1916}
Function	ATP-dependent chromatin remodeler that mediates chromatin-remodeling following DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 33357431 , PubMed: 34210977 , PubMed: 34486521 , PubMed: 34874266). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653). Chromatin remodeling activity promotes PARP2 removal from chromatin (PubMed: 33275888).
Cellular Location	Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by

histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.


Tissue Location

Frequently overexpressed in hepatomacellar carcinomas.

Background

DNA helicase which plays a role in chromatin-remodeling following DNA damage. Targeted to sites of DNA damage through interaction with poly(ADP-ribose) and functions to regulate chromatin during DNA repair. Able to catalyze nucleosome sliding in an ATP-dependent manner. Helicase activity is strongly stimulated upon poly(ADP-ribose)-binding.

Images



Western Blot analysis of extracts from NIH-3T3, Jurkat, A549 cells, using CHD1L Polyclonal Antibody.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.