

TudorSN Polyclonal Antibody

Catalog # AP73267

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q7KZF4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101997

Additional Information

Gene ID	27044
Other Names	SND1; TDRD11; Staphylococcal nuclease domain-containing protein 1; 100 kDa coactivator; EBNA2 coactivator p100; Tudor domain-containing protein 11; p100 co-activator
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SND1
Synonyms	TDRD11
Function	Endonuclease that mediates miRNA decay of both protein-free and AGO2-loaded miRNAs (PubMed: 18453631 , PubMed: 28546213). As part of its function in miRNA decay, regulates mRNAs involved in G1-to-S phase transition (PubMed: 28546213). Functions as a bridging factor between STAT6 and the basal transcription factor (PubMed: 12234934). Plays a role in PIM1 regulation of MYB activity (PubMed: 9809063). Functions as a transcriptional coactivator for STAT5 (By similarity).
Cellular Location	Cytoplasm. Nucleus. Melanosome Note=In IL-4 stimulated cells colocalizes with STAT6 in the nucleus (PubMed:12234934). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065)


Tissue Location

Ubiquitously expressed.


Background

Endonuclease that mediates miRNA decay of both protein-free and AGO2-loaded miRNAs (PubMed:[28546213](#), PubMed:[18453631](#)). As part of its function in miRNA decay, regulates mRNAs involved in G1-to-S phase transition (PubMed:[28546213](#)). Functions as a bridging factor between STAT6 and the basal transcription factor (PubMed:[12234934](#)). Plays a role in PIM1 regulation of MYB activity (PubMed:[9809063](#)). Functions as a transcriptional coactivator for STAT5 (By similarity).


Images




Western Blot analysis of extracts from Jurkat cells, using TudorSN Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.