

PAOX Polyclonal Antibody

Catalog # AP73281

Product Information

Application	WB, IHC-P
Primary Accession	Q6QHF9
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55513

Additional Information

Gene ID	196743
Other Names	PAOX; PAO; Peroxisomal N(1)-acetyl-spermine/spermidine oxidase; Polyamine oxidase
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	PAOX
Synonyms	PAO
Function	Flavoenzyme which catalyzes the oxidation of N(1)- acetylspermine to spermidine and is thus involved in the polyamine back-conversion (PubMed: 12477380). Can also oxidize N(1)- acetylspermidine to putrescine. Substrate specificity: N(1)- acetylspermine = N(1)-acetylspermidine > N(1),N(12)-diacetylspermine >> spermine. Does not oxidize spermidine. Plays an important role in the regulation of polyamine intracellular concentration and has the potential to act as a determinant of cellular sensitivity to the antitumor polyamine analogs (PubMed: 12477380).
Cellular Location	Peroxisome. Cytoplasm
Tissue Location	Widely expressed. Not detected in spleen. Expressed at lower level in neoplastic tissues.


Background

Flavoenzyme which catalyzes the oxidation of N(1)-acetylspermine to spermidine and is thus involved in the polyamine back-conversion. Can also oxidize N(1)-acetylspermidine to putrescine. Substrate specificity: N(1)-acetylspermine = N(1)-acetylspermidine > N(1),N(12)-diacetylspermine >> spermine. Does not oxidize spermidine. Plays an important role in the regulation of polyamine intracellular concentration and has the potential to act as a determinant of cellular sensitivity to the antitumor polyamine analogs.


Images



Western Blot analysis of extracts from rat stomach, K562 cells, using PAOX Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.