

SLC6A14 Polyclonal Antibody

Catalog # AP73285

Product Information

Application	WB, IHC-P
Primary Accession	Q9UN76
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	72153

Additional Information

Gene ID	11254
Other Names	SLC6A14; Sodium- and chloride-dependent neutral and basic amino acid transporter B(0+; Amino acid transporter ATB0+; Solute carrier family 6 member 14
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	SLC6A14 (HGNC:11047)
Function	Amino acid transporter that plays an important role in the absorption of amino acids in the intestinal tract. Mediates the uptake of a broad range of neutral and cationic amino acids (with the exception of proline) in a Na(+)/Cl(-)-dependent manner (PubMed: 10446133). Transports non-alpha-amino acids such as beta- alanine with low affinity, and has a higher affinity for dipolar and cationic amino acids such as leucine and lysine (PubMed: 18599538). Can also transport carnitine, butyrylcarnitine and propionylcarnitine coupled to the transmembrane gradients of Na(+) and Cl(-) (PubMed: 17855766).
Cellular Location	Membrane; Multi- pass membrane protein. Apical cell membrane {ECO:0000250 UniProtKB:Q9JMA9}; Multi-pass membrane protein
Tissue Location	Levels are highest in adult and fetal lung, in trachea and salivary gland. Lower levels detected in mammary gland, stomach and pituitary gland, and very low

levels in colon, uterus, prostate and testis.


Background

Mediates the uptake of a broad range of neutral and cationic amino acids (with the exception of proline) in a Na(+)/Cl(-)-dependent manner.

Images



Western Blot analysis of extracts from 293 cells, using SLC6A14 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.