

SLC6A14 Polyclonal Antibody

Catalog # AP73285

Product Information

Application WB, IHC-P
Primary Accession Q9UN76
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 72153

Additional Information

Gene ID 11254

Other Names SLC6A14; Sodium- and chloride-dependent neutral and basic amino acid

transporter B(0+; Amino acid transporter ATB0+; Solute carrier family 6

member 14

Dilution WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet

tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name SLC6A14 (HGNC:11047)

Function Amino acid transporter that plays an important role in the absorption of

amino acids in the intestinal tract. Mediates the uptake of a broad range of neutral and cationic amino acids (with the exception of proline) in a

Na(+)/Cl(-)-dependent manner (PubMed: 10446133). Transports non-alpha-amino acids such as beta- alanine with low affinity, and has a

higher affinity for dipolar and cationic amino acids such as leucine and lysine (PubMed: 18599538). Can also transport carnitine, butirylcarnitine and propionylcarnitine coupled to the transmembrane gradients of Na(+) and Cl(-)

(PubMed: 17855766).

Cellular Location Membrane; Multi- pass membrane protein. Apical cell membrane

{ECO:0000250|UniProtKB:Q9JMA9}; Multi-pass membrane protein

Tissue Location Levels are highest in adult and fetal lung, in trachea and salivary gland. Lower


levels detected in mammary gland, stomach and pituitary gland, and very low

levels in colon, uterus, prostate and testis.


Background

Mediates the uptake of a broad range of neutral and cationic amino acids (with the exception of proline) in a Na(+)/Cl(-)-dependent manner.

Images



Western Blot analysis of extracts from 293 cells, using SLC6A14 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.