

NCX1 Polyclonal Antibody

Catalog # AP73286

Product Information

Application	WB
Primary Accession	P32418
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	108547

Additional Information

Gene ID	6546
Other Names	SLC8A1; CNC; NCX1; Sodium/calcium exchanger 1; Na(+)/Ca(2+)-exchange protein 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	SLC8A1
Function	Mediates the exchange of one Ca(2+) ion against three to four Na(+) ions across the cell membrane, and thereby contributes to the regulation of cytoplasmic Ca(2+) levels and Ca(2+)-dependent cellular processes (PubMed: 11241183 , PubMed: 1374913 , PubMed: 1476165). Contributes to Ca(2+) transport during excitation-contraction coupling in muscle (PubMed: 11241183 , PubMed: 1374913 , PubMed: 1476165). In a first phase, voltage-gated channels mediate the rapid increase of cytoplasmic Ca(2+) levels due to release of Ca(2+) stores from the endoplasmic reticulum (PubMed: 11241183 , PubMed: 1374913 , PubMed: 1476165). SLC8A1 mediates the export of Ca(2+) from the cell during the next phase, so that cytoplasmic Ca(2+) levels rapidly return to baseline (PubMed: 11241183 , PubMed: 1374913 , PubMed: 1476165). Required for normal embryonic heart development and the onset of heart contractions (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Detected primarily in heart and at lower levels in brain (PubMed: 1374913).

Expressed in cardiac sarcolemma, brain, kidney, liver, pancreas, skeletal muscle, placenta and lung (PubMed:1476165)

Background

Mediates the exchange of one Ca^{2+} ion against three to four Na^{+} ions across the cell membrane, and thereby contributes to the regulation of cytoplasmic Ca^{2+} levels and Ca^{2+} - dependent cellular processes (PubMed:[1374913](#), PubMed:[11241183](#), PubMed:[1476165](#)). Contributes to Ca^{2+} transport during excitation-contraction coupling in muscle. In a first phase, voltage-gated channels mediate the rapid increase of cytoplasmic Ca^{2+} levels due to release of Ca^{2+} stores from the endoplasmic reticulum. SLC8A1 mediates the export of Ca^{2+} from the cell during the next phase, so that cytoplasmic Ca^{2+} levels rapidly return to baseline. Required for normal embryonic heart development and the onset of heart contractions.

Images



Western Blot analysis of extracts from 293 cells, using NCX1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.