

SCP-3 Polyclonal Antibody

Catalog # AP73300

Product Information

Application	WB
Primary Accession	Q8IZU3
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27729

Additional Information

Gene ID	50511
Other Names	SYCP3; SCP3; Synaptonemal complex protein 3; SCP-3
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	SYCP3
Synonyms	SCP3
Function	Component of the synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase. Required for centromere pairing during meiosis in male germ cells (By similarity). Required for normal meiosis during spermatogenesis and male fertility (PubMed: 14643120). Plays a lesser role in female fertility. Required for efficient phosphorylation of HORMAD1 and HORMAD2 (By similarity).
Cellular Location	Nucleus {ECO:0000250 UniProtKB:Q60547}. Chromosome {ECO:0000250 UniProtKB:Q60547}. Chromosome, centromere {ECO:0000250 UniProtKB:Q60547}. Note=It is present in early unpaired cores, in the lateral domains of the synaptonemal complex and in the chromosome cores when they separate at diplotene. It is found axial to the metaphase I chromosomes and in association with pairs of sister centromeres. The centromere-associated protein becomes dissociated from the centromeres at anaphase II and is not found in mitotic metaphase centromeres. {ECO:0000250 UniProtKB:Q60547}

Background

Component of the synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase. Required for centromere pairing during meiosis in male germ cells (By similarity). Required for normal meiosis during spermatogenesis and male fertility (PubMed:[14643120](#)). Plays a lesser role in female fertility. Required for efficient phosphorylation of HORMAD1 and HORMAD2 (By similarity).

Images



Western Blot analysis of K562, 22RV-1, H460 cells using SCP-3 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.