

MMP-12 Polyclonal Antibody

Catalog # AP73320

Product Information

Application WB, IHC-P **Primary Accession** P39900

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW54002

Additional Information

Gene ID 4321

Other Names MMP12; HME; Macrophage metalloelastase; MME; Macrophage elastase; ME;

hME; Matrix metalloproteinase-12; MMP-12

Dilution WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet

tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p:

1:100-300 ELISA: 1/20000. Not yet tested in other applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name MMP12

Synonyms HME

Function May be involved in tissue injury and remodeling. Has significant elastolytic

activity. Can accept large and small amino acids at the P1' site, but has a preference for leucine. Aromatic or hydrophobic residues are preferred at the P1 site, with small hydrophobic residues (preferably alanine) occupying P3.

Cellular Location Secreted, extracellular space, extracellular matrix


Tissue Location Found in alveolar macrophages but not in peripheral blood monocytes

Background


May be involved in tissue injury and remodeling. Has significant elastolytic activity. Can accept large and small amino acids at the P1' site, but has a preference for leucine. Aromatic or hydrophobic residues are

preferred at the P1 site, with small hydrophobic residues (preferably alanine) occupying P3.

Images



Western Blot analysis of Jurkat cells using MMP-12 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.