Sg III Polyclonal Antibody

Catalog # AP73325

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q8WXD2
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53005

Additional Information

|	
Gene ID	29106
Other Names	SCG3; Secretogranin-3; Secretogranin III; SgIII
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information

|	
Name	SCG3
Function	Member of the granin protein family that regulates the biogenesis of secretory granules (PubMed:19357184). Acts as a sorting receptor for intragranular proteins including chromogranin A/CHGA (By similarity). May also play a role in angiogenesis. Promotes endothelial proliferation, migration and tube formation through MEK/ERK signaling pathway (PubMed:29154827).
Cellular Location	Cytoplasmic vesicle, secretory vesicle {ECO:0000250|UniProtKB:P47868}. Cytoplasmic vesicle, secretory vesicle membrane {ECO:0000250|UniProtKB:P47868}; Peripheral membrane protein. Secreted. Note=Associated with the secretory granule membrane through direct binding to cholesterol-enriched lipid rafts. {ECO:0000250|UniProtKB:P47868}
Tissue Location	Detected in urine (at protein level) (PubMed:25326458). Expressed in brain, heart, kidney, liver and skeletal muscle.

Images

Western Blot analysis of mouse heart, mouse kidney cells using Sg III Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.