

Sg III Polyclonal Antibody

Catalog # AP73325

Product Information

Application WB, IHC-P
Primary Accession Q8WXD2
Reactivity Human, Rat
Host Rabbit
Clonality Polyclonal
Calculated MW 53005

Additional Information

Gene ID 29106

Other Names SCG3; Secretogranin-3; Secretogranin III; SgIII

Dilution WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet

tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p:

1:100-300 ELISA: 1/20000. Not yet tested in other applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name SCG3

Function Member of the granin protein family that regulates the biogenesis of

secretory granules (PubMed: 19357184). Acts as a sorting receptor for intragranular proteins including chromogranin A/CHGA (By similarity). May also play a role in angiogenesis. Promotes endothelial proliferation, migration and tube formation through MEK/ERK signaling pathway (PubMed: 29154827).

Cellular Location Cytoplasmic vesicle, secretory vesicle {ECO:0000250 | UniProtKB:P47868}.


Cytoplasmic vesicle, secretory vesicle membrane

{ECO:0000250 | UniProtKB:P47868}; Peripheral membrane protein. Secreted. Note=Associated with the secretory granule membrane through direct binding to cholesterol-enriched lipid rafts. {ECO:0000250 | UniProtKB:P47868}


Tissue Location Detected in urine (at protein level) (PubMed:25326458). Expressed in brain,

heart, kidney, liver and skeletal muscle.


Images



Western Blot analysis of mouse heart, mouse kidney cells using Sg III Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.