

# CYP11A1 Polyclonal Antibody

Catalog # AP73406

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">P05108</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60102

## Additional Information

Gene ID	1583
Other Names	CYP11A1; CYP11A; Cholesterol side-chain cleavage enzyme, mitochondrial; CYPXIA1; Cholesterol desmolase; Cytochrome P450 11A1; Cytochrome P450(scc)
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


## Protein Information

Name	CYP11A1 {ECO:0000303   PubMed:21636783, ECO:0000312   HGNC:HGNC:2590}
Function	A cytochrome P450 monooxygenase that catalyzes the side-chain hydroxylation and cleavage of cholesterol to pregnenolone, the precursor of most steroid hormones (PubMed: <a href="#">21636783</a> ). Catalyzes three sequential oxidation reactions of cholesterol, namely the hydroxylation at C22 followed with the hydroxylation at C20 to yield 20R,22R- hydroxycholesterol that is further cleaved between C20 and C22 to yield the C21-steroid pregnenolone and 4-methylpentanal (PubMed: <a href="#">21636783</a> ). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate and reducing the second into a water molecule. Two electrons are provided by NADPH via a two-protein mitochondrial transfer system comprising flavoprotein FDXR (adrenodoxin/ferredoxin reductase) and nonheme iron-sulfur protein FDX1 or FDX2 (adrenodoxin/ferredoxin) (PubMed: <a href="#">21636783</a> ).
Cellular Location	Mitochondrion inner membrane {ECO:0000250   UniProtKB:P14137}; Peripheral membrane protein. Note=Localizes to the matrix side of the


## Background

Catalyzes the side-chain cleavage reaction of cholesterol to pregnenolone, the precursor of most steroid hormones.


## Images




Western Blot analysis of K562 cells using CYP11A1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100