

# SPTLC1 Polyclonal Antibody

Catalog # AP73415

## **Product Information**

Application	WB, IHC-P
Primary Accession	<u>015269</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52744

#### **Additional Information**

Gene ID	10558
Other Names	SPTLC1; LCB1; Serine palmitoyltransferase 1; Long chain base biosynthesis protein 1; LCB 1; Serine-palmitoyl-CoA transferase 1; SPT 1; SPT1
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

### **Protein Information**

Name	SPTLC1
Synonyms	LCB1
Function	Component of the serine palmitoyltransferase multisubunit enzyme (SPT) that catalyzes the initial and rate-limiting step in sphingolipid biosynthesis by condensing L-serine and activated acyl-CoA (most commonly palmitoyl-CoA) to form long-chain bases. The SPT complex is also composed of SPTLC2 or SPTLC3 and SPTSSA or SPTSSB. Within this complex, the heterodimer with SPTLC2 or SPTLC3 forms the catalytic core (PubMed: <u>19416851</u> , PubMed: <u>33558762</u> , PubMed: <u>33558762</u> ). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed: <u>19416851</u> , PubMed: <u>33558762</u> ). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1-SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed: <u>19416851</u> , PubMed: <u>19648650</u> ). The SPTLC1-SPTLC3-SPTSSB isozyme

	displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed: <u>19416851</u> , PubMed: <u>19648650</u> , PubMed: <u>33558761</u> , PubMed: <u>33558762</u> ). Required for adipocyte cell viability and metabolic homeostasis (By similarity).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein {ECO:0000250 UniProtKB:O35704}
Tissue Location	Widely expressed. Not detected in small intestine.

# Background

Serine palmitoyltransferase (SPT) (PubMed: <u>19416851</u>). The heterodimer formed with SPTLC2 or SPTLC3 constitutes the catalytic core (PubMed:<u>19416851</u>). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed:<u>19416851</u>). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1- SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed:<u>19416851</u>). The SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC3-SPTSSB isozyme displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed:<u>19416851</u>). Required for adipocyte cell viability and metabolic homeostasis (By similarity).

#### Images



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.