

# DNAM-1 Polyclonal Antibody

Catalog # AP73585

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q15762</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38614

## Additional Information

---

Gene ID	10666
Other Names	CD226; DNAM1; CD226 antigen; DNAX accessory molecule 1; DNAM-1; CD226
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	CD226
Synonyms	DNAM1
Function	<p>Cell surface receptor that plays an important role in the immune system, particularly in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-cells and NK cells (PubMed:<a href="#">8673704</a>, PubMed:<a href="#">9712030</a>). Functions as a costimulatory receptor upon recognition of target cells, such as virus- infected or tumor cells. Upon binding to its ligands PVR/CD155 or NECTIN2/CD112 on target cells, promotes the cytotoxic activity of NK cells and CTLs, enhancing their ability to kill these cells (PubMed:<a href="#">26755705</a>, PubMed:<a href="#">31253644</a>, PubMed:<a href="#">30591568</a>).</p> <p>Mechanistically, phosphorylation by Src kinases such as LYN of FYN, enables binding to adapter GRB2, leading to activation of VAV1, PI3K and PLCG1. Promotes also activation of kinases ERK and AKT, as well as calcium fluxes (By similarity).</p>
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Localizes to lipid rafts.

**Tissue Location**

Expressed by peripheral blood T-lymphocytes.


**Background**

---

Involved in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-lymphocyte (CTL) and NK cell (PubMed:[8673704](#)). Cell surface receptor for NECTIN2. Upon ligand binding, stimulates T- cell proliferation and cytokine production, including that of IL2, IL5, IL10, IL13, and IFNG. Competes with PVRIG for NECTIN2-binding (PubMed:[26755705](#)).

**Images**

---



Western Blot analysis of HeLa cells using DNAM-1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.