

DCL-1 Polyclonal Antibody

Catalog # AP73656

Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q8IX05 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 26183 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 9936 |
| Other Names | CD302; CLEC13A; DCL1; KIAA0022; CD302 antigen; C-type lectin BIMLEC; C-type lectin domain family 13 member A; DEC205-associated C-type lectin 1; Type I transmembrane C-type lectin receptor DCL-1; CD302 |
| Dilution | WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |

Protein Information

| | |
|-------------------|--|
| Name | CD302 |
| Synonyms | CLEC13A, DCL1, KIAA0022 |
| Function | Potential multifunctional C-type lectin receptor that may play roles in endocytosis and phagocytosis as well as in cell adhesion and migration. |
| Cellular Location | Membrane; Single-pass type I membrane protein. Cell projection, filopodium. Cytoplasm, cell cortex. Cell projection, microvillus. Note=Colocalizes with F-actin in filopodia, cellular cortex and microvilli of the apical cell surface |
| Tissue Location | Expressed at moderate levels in monocytes, myeloid blood dendritic cells and granulocytes and at low levels in plasmacytoid blood dendritic cells, monocyte-derived macrophages and monocyte-derived dendritic cells, with no expression detected in T- lymphocytes, B-lymphocytes and natural killer cells (at protein level) Expressed widely in different tissues, with highest expression levels in liver, lung, peripheral blood leukocytes and spleen, and lowest levels in neuronal tissues, skeletal muscle and ovary. Isoform 2 and |

isoform 3 are expressed in malignant Hodgkin lymphoma cells called Hodgkin and Reed-Sternberg (HRS) cells.

Background

Potential multifunctional C-type lectin receptor that may play roles in endocytosis and phagocytosis as well as in cell adhesion and migration.

Images



Western Blot analysis of HeLa cells using DCL-1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.