

TRAIL Polyclonal Antibody

Catalog # AP73694

Product Information

Application	WB, IHC-P
Primary Accession	P50591
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32509

Additional Information

Gene ID	8743
Other Names	TNFSF10; APO2L; TRAIL; Tumor necrosis factor ligand superfamily member 10; Apo-2 ligand; Apo-2L; TNF-related apoptosis-inducing ligand; Protein TRAIL; CD253
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	TNFSF10
Synonyms	APO2L, TRAIL
Function	Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG (PubMed: 10549288 , PubMed: 26457518). Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis.
Cellular Location	Cell membrane; Single-pass type II membrane protein. Secreted. Note=Exists both as membrane-bound and soluble form.
Tissue Location	Widespread; most predominant in spleen, lung and prostate


Background

Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG (PubMed:[26457518](#), PubMed:[10549288](#)). Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis.


Images




Western Blot analysis of HeLa cells using TRAIL Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human lymph nodes, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human tonsilla, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human tonsilla, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.