

OAS2 Polyclonal Antibody

Catalog # AP73719

Product Information

Application	WB, IHC-P
Primary Accession	P29728
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82431

Additional Information

Gene ID	4939
Other Names	OAS2; 2'-5'-oligoadenylate synthase 2; (2-5')oligo(A) synthase 2; 2-5A synthase 2; p69 OAS / p71 OAS; p69OAS / p71OAS
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	OAS2 (HGNC:8087)
Function	Interferon-induced, dsRNA-activated antiviral enzyme which plays a critical role in cellular innate antiviral response (PubMed: 10464285 , PubMed: 9880569). Activated by detection of double stranded RNA (dsRNA): polymerizes higher oligomers of 2'-5'- oligoadenylates (2-5A) from ATP which then bind to the inactive monomeric form of ribonuclease L (RNASEL) leading to its dimerization and subsequent activation (PubMed: 10464285 , PubMed: 11682059 , PubMed: 9880569). Activation of RNASEL leads to degradation of cellular as well as viral RNA, resulting in the inhibition of protein synthesis, thus terminating viral replication (PubMed: 10464285 , PubMed: 9880569). Can mediate the antiviral effect via the classical RNASEL-dependent pathway or an alternative antiviral pathway independent of RNASEL (PubMed: 21142819). In addition, it may also play a role in other cellular processes such as apoptosis, cell growth, differentiation and gene regulation (PubMed: 21142819). May act as a negative regulator of lactation, stopping lactation in virally infected mammary gland lobules, thereby preventing transmission of viruses to neonates (By similarity). Non-infected

lobules would not be affected, allowing efficient pup feeding during infection (By similarity).


Cellular Location

Cytoplasm. Cytoplasm, perinuclear region


Background

Interferon-induced, dsRNA-activated antiviral enzyme which plays a critical role in cellular innate antiviral response (PubMed:[10464285](#), PubMed:[9880569](#)). Activated by detection of double stranded RNA (dsRNA): polymerizes higher oligomers of 2'- 5'-oligoadenylates (2-5A) from ATP which then bind to the inactive monomeric form of ribonuclease L (RNASEL) leading to its dimerization and subsequent activation (PubMed:[10464285](#), PubMed:[9880569](#), PubMed:[11682059](#)). Activation of RNASEL leads to degradation of cellular as well as viral RNA, resulting in the inhibition of protein synthesis, thus terminating viral replication (PubMed:[10464285](#), PubMed:[9880569](#)). Can mediate the antiviral effect via the classical RNASEL-dependent pathway or an alternative antiviral pathway independent of RNASEL (PubMed:[21142819](#)). In addition, it may also play a role in other cellular processes such as apoptosis, cell growth, differentiation and gene regulation (PubMed:[21142819](#)). May act as a negative regulator of lactation, stopping lactation in virally infected mammary gland lobules, thereby preventing transmission of viruses to neonates (By similarity). Non-infected lobules would not be affected, allowing efficient pup feeding during infection (By similarity).


Images



Western Blot analysis of HeLa cells using OAS2 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-uterus, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-uterus, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.