

ApoC-III Polyclonal Antibody

Catalog # AP73733

Product Information

Application	WB, IHC-P
Primary Accession	P02656
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10852

Additional Information

Gene ID	345
Other Names	APOC3; Apolipoprotein C-III; Apo-CIII; ApoC-III; Apolipoprotein C3
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	APOC3
Function	Component of triglyceride-rich very low density lipoproteins (VLDL) and high density lipoproteins (HDL) in plasma (PubMed: 18201179 , PubMed: 22510806). Plays a multifaceted role in triglyceride homeostasis (PubMed: 18201179 , PubMed: 22510806). Intracellularly, promotes hepatic very low density lipoprotein 1 (VLDL1) assembly and secretion; extracellularly, attenuates hydrolysis and clearance of triglyceride- rich lipoproteins (TRLs) (PubMed: 18201179 , PubMed: 22510806). Impairs the lipolysis of TRLs by inhibiting lipoprotein lipase and the hepatic uptake of TRLs by remnant receptors (PubMed: 18201179 , PubMed: 22510806). Formed of several curved helices connected via semiflexible hinges, so that it can wrap tightly around the curved micelle surface and easily adapt to the different diameters of its natural binding partners (PubMed: 18408013).
Cellular Location	Secreted
Tissue Location	Liver..


Background

Component of triglyceride-rich very low density lipoproteins (VLDL) and high density lipoproteins (HDL) in plasma (PubMed:[18201179](#), PubMed:[22510806](#)). Plays a multifaceted role in triglyceride homeostasis (PubMed:[18201179](#), PubMed:[22510806](#)). Intracellularly, promotes hepatic very low density lipoprotein 1 (VLDL1) assembly and secretion; extracellularly, attenuates hydrolysis and clearance of triglyceride-rich lipoproteins (TRLs) (PubMed:[18201179](#), PubMed:[22510806](#)). Impairs the lipolysis of TRLs by inhibiting lipoprotein lipase and the hepatic uptake of TRLs by remnant receptors (PubMed:[18201179](#), PubMed:[22510806](#)). Formed of several curved helices connected via semiflexible hinges, so that it can wrap tightly around the curved micelle surface and easily adapt to the different diameters of its natural binding partners (PubMed:[18408013](#)).

Images



Western Blot analysis of 293T-UV cells using ApoC-III Polyclonal Antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.