10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD53 Polyclonal Antibody

Catalog # AP73774

Product Information

Application	WB, E
Primary Accession	P19397
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24341

Additional Information

Gene ID	963
Other Names	CD53; MOX44; TSPAN25; Leukocyte surface antigen CD53; Cell surface glycoprotein CD53; Tetraspanin-25; Tspan-25; CD53
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	CD53
Synonyms	MOX44, TSPAN25
Function	Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling (PubMed:28487417). Participates thereby in diverse biological functions such as cell signal transduction, adhesion, migration and protein trafficking (PubMed:32974937, PubMed:35767951). Plays a role in the activation of monocytes and B-cells (PubMed:8335905). Acts as an essential regulator of B-cell development by promoting interleukin-7 receptor/IL7R signaling (By similarity). Also promotes, in B-cells, the BCR signaling by recruiting PKC to the plasma membrane in order to phosphorylate its substrates (PubMed:28487417). Plays an essential role in B- and T-cells homing to lymph nodes by stabilizing L-selectin/SELL cell surface expression (By similarity). Also mediates metabolic and inflammatory functions in hepatocytes and adipose tissue by promoting TNF and LPS signaling independent of the immune compartment (By similarity).

Cellular Location	Cell membrane. Cell junction {ECO:0000250|UniProtKB:Q61451}. Membrane; Multi-pass membrane protein. Synapse. Note=Concentrates in localized microdomains along the plasma membrane at the contact sites between cells of fused myotubes. {ECO:0000250|UniProtKB:Q61451}
Tissue Location	B-cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive thymocytes and peripheral T-cells


Background

Required for efficient formation of myofibers in regenerating muscle at the level of cell fusion. May be involved in growth regulation in hematopoietic cells (By similarity).

Images



Western Blot analysis of KB cells using CD53 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of KB cells using CD53 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.