

Blr1 Polyclonal Antibody

Catalog # AP73796

Product Information

Application	WB
Primary Accession	P32302
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41955

Additional Information

Gene ID	643
Other Names	CXCR5; BLR1; MDR15; C-X-C chemokine receptor type 5; CXC-R5; CXCR-5; Burkitt lymphoma receptor 1; Monocyte-derived receptor 15; MDR-15; CD185
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	CXCR5
Synonyms	BLR1, MDR15
Function	Cytokine receptor that binds to B-lymphocyte chemoattractant (BLC). Involved in B-cell migration into B-cell follicles of spleen and Peyer patches but not into those of mesenteric or peripheral lymph nodes. May have a regulatory function in Burkitt lymphoma (BL) lymphomagenesis and/or B-cell differentiation.
Cellular Location	Cell membrane; Multi-pass membrane protein.
Tissue Location	Expression in mature B-cells and Burkitt lymphoma cells

Background


Cytokine receptor that binds to B-lymphocyte chemoattractant (BLC). Involved in B-cell migration into B-cell follicles of spleen and Peyer patches but not into those of mesenteric or peripheral lymph nodes. May

have a regulatory function in Burkitt lymphoma (BL) lymphomagenesis and/or B-cell differentiation.

Images



Western Blot analysis of NIH-3T3 cells using Blr1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of 3T3 cells using Blr1 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.