

CD192 Polyclonal Antibody

Catalog # AP73818

Product Information

Application	WB
Primary Accession	P41597
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41915

Additional Information

Gene ID	729230
Other Names	CCR2; CMKBR2; C-C chemokine receptor type 2; C-C CKR-2; CC-CKR-2; CCR-2; CCR2; Monocyte chemoattractant protein 1 receptor; MCP-1-R; CD192
Dilution	WB--Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CCR2
Synonyms	CMKBR2
Function	Key functional receptor for CCL2 but can also bind CCL7, and CCL12 (PubMed: 23408426 , PubMed: 38157855 , PubMed: 8048929 , PubMed: 8146186). Also transduces signaling mediated by CCL13 (PubMed: 38157855). Its binding with CCL2 on monocytes and macrophages mediates chemotaxis and migration induction through the activation of the PI3K cascade, the small G protein Rac and lamellipodium protrusion (PubMed: 38157855). Also acts as a receptor for the beta-defensin DEFB106A/DEFB106B (PubMed: 23938203). Regulates the expression of T-cell inflammatory cytokines and T-cell differentiation, promoting the differentiation of T-cells into T-helper 17 cells (Th17) during inflammation (By similarity). Facilitates the export of mature thymocytes by enhancing directional movement of thymocytes to sphingosine-1-phosphate stimulation and up-regulation of S1P1R expression; signals through the JAK-STAT pathway to regulate FOXO1 activity leading to an increased expression of S1P1R (By similarity). Plays an important role in mediating peripheral nerve injury-induced neuropathic pain (By similarity).

Increases NMDA-mediated synaptic transmission in both dopamine D1 and D2 receptor-containing neurons, which may be caused by MAPK/ERK-dependent phosphorylation of GRIN2B/NMDAR2B (By similarity). Mediates the recruitment of macrophages and monocytes to the injury site following brain injury (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Note=The chemoattractant receptors are distributed throughout the cell surface; after stimulation with a ligand, such as CCL2, they are rapidly recruited into microdomain clusters at the cell membrane.


Tissue Location

Expressed by monocytes and IL2-activated NK cells (PubMed:9058802). Abundantly expressed on CD14+/CD16- monocytes and weakly on CD14+/CD16+ monocytes, type 2 dendritic cells (DCs) and plasmacytoid DCs (at protein level) (PubMed:38157855)

Background


Key functional receptor for CCL2 but can also bind CCL7 and CCL12 (PubMed: [8146186](#), PubMed:[8048929](#), PubMed:[23408426](#)). Its binding with CCL2 on monocytes and macrophages mediates chemotaxis and migration induction through the activation of the PI3K cascade, the small G protein Rac and lamellipodium protrusion (Probable). Also acts as a receptor for the beta-defensin DEFB106A/DEFB106B (PubMed:[23938203](#)). Regulates the expression of T-cell inflammatory cytokines and T-cell differentiation, promoting the differentiation of T-cells into T-helper 17 cells (Th17) during inflammation (By similarity). Facilitates the export of mature thymocytes by enhancing directional movement of thymocytes to sphingosine-1-phosphate stimulation and up- regulation of S1P1R expression; signals through the JAK-STAT pathway to regulate FOXO1 activity leading to an increased expression of S1P1R (By similarity). Plays an important role in mediating peripheral nerve injury-induced neuropathic pain (By similarity). Increases NMDA-mediated synaptic transmission in both dopamine D1 and D2 receptor-containing neurons, which may be caused by MAPK/ERK-dependent phosphorylation of GRIN2B/NMDAR2B (By similarity). Mediates the recruitment of macrophages and monocytes to the injury site following brain injury (By similarity).

Images



Western Blot analysis of NIH-3T3 cells using CD192 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Western Blot analysis of 3T3 cells using CD192 Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.