

C7 Polyclonal Antibody

Catalog # AP73883

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P10643
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	93518

Additional Information

Gene ID	730
Other Names	C7; Complement component C7
Dilution	WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	C7 {ECO:0000303 PubMed:3335508, ECO:0000312 HGNC:HGNC:1346}
Function	Component of the membrane attack complex (MAC), a multiprotein complex activated by the complement cascade, which inserts into a target cell membrane and forms a pore, leading to target cell membrane rupture and cell lysis (PubMed: 22832194 , PubMed: 26841837 , PubMed: 27052168 , PubMed: 30552328 , PubMed: 3335508). The MAC is initiated by proteolytic cleavage of C5 into complement C5b in response to the classical, alternative, lectin and GZMK complement pathways (PubMed: 22832194 , PubMed: 30552328 , PubMed: 3335508). The complement pathways consist in a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed: 22832194 , PubMed: 30552328 , PubMed: 3335508). C7 serves as a membrane anchor (PubMed: 30552328). During MAC assembly, associates with C5b and C6 to form the C5b-7 complex, a key lipophilic precursor of the MAC complex, which associates with the outer leaflet and reduces the energy for membrane bending (PubMed: 30552328 , PubMed: 32569291).


Cellular Location

Secreted. Target cell membrane Note=Secreted as soluble protein (PubMed:3335508). Inserts into the cell membrane of target cells (PubMed:30552328, PubMed:31061395)


Background

Constituent of the membrane attack complex (MAC) that plays a key role in the innate and adaptive immune response by forming pores in the plasma membrane of target cells. C7 serves as a membrane anchor.


Images



Western blot analysis of Hela lysis using C7 antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.